

## Evaluation of Antioxidant Capacities and total Polyphenols in Various Edible Parts of *Capparis spinosa* L. Collected from trans-Himalayas

Manish S. Bhojar<sup>#</sup>, Gyan P. Mishra<sup>#,\*</sup>, Pradeep K. Naik<sup>1</sup>, and Shashi Bala Singh<sup>#</sup>

<sup>#</sup>DRDO-Defence Institute of High Altitude Research, Leh-194101, India

<sup>1</sup>Jaypee University of Information Technology, Wanknaghat, Solan-173234, India

\*E-mail: gyan.gene@gmail.com

### ABSTRACT

The phytochemical screening, antioxidant capacity, and total polyphenols in the methanolic extract of leaves, flower buds, roots and fruits of *Capparis spinosa* collected from trans-Himalayan region of Ladakh were assessed in an effort to corroborate its medicinal and culinary potential. Highest DPPH and ABTS radical scavenging activity were observed in the leaves and least in dried fruit samples, even FRAP assay also illustrated the same trend. IC50 values of DPPH assay was highly correlated with that of ABTS ( $R^2 = 0.9084$ ) and FRAP assay ( $R^2 = 0.9771$ ). However, IC50 value of ABTS was reasonably correlated with FRAP assay ( $R^2 = 0.5838$ ). The highest phenolic and flavonoid content was recorded in the leaf samples (24.78 and 5.69 mg GAE/g DW respectively), whereas it was lowest in the dried fruit samples (4.07 mg quercetin equivalent/g DW and nil, respectively). The total phenolic contents were highly correlated with IC50 value of ABTS ( $R^2 = 0.9084$ ), DPPH ( $R^2 = 0.9388$ ) and FRAP value ( $R^2 = 0.9618$ ). But, total flavonoid contents were highly correlated with ABTS ( $R^2 = 0.7449$ ), DPPH ( $R^2 = 0.8791$ ) and FRAP values ( $R^2 = 0.9588$ ). Thus, this study has validated the medicinal potential of all the edible parts of the *C. spinosa*.

**Keywords:** Capers; Phytochemical screening; Antioxidant activity; Polyphenols; Correlation

### 1. INTRODUCTION

There is a growing demand for natural products in the human diet, both due to the possible negative effects of synthetic food additives on human health and also to increase consumer perception of this problem in recent years. The edible parts of *Capparis* like flower buds, fruits, leaves, and root barks are known to contain a range of antioxidants, including polyphenolic compounds, which are significantly correlated with antioxidant potential<sup>1,2</sup>.

*Capparis spinosa* L. (Capparidaceae) also called 'Caper' and locally known as 'Kabra' is one of the oldest known medicinal plants in 'Amchi system' which is local medicinal system of Ladakh (India). It has been utilised in preparations of various herbal formulations since long time for the treatment of a range of ailments like gastrointestinal infection, diarrhea, diabetes, rheumatism and also used as a leafy vegetable and forage by local people residing in trans-Himalayan region of Ladakh<sup>3,4</sup>. In India, it is found in the inner valleys of trans-Himalayas between 3020 m – 3890 m above mean sea level (AMSL) which includes Indus, Nubra and Suru valleys of Ladakh region. It is an under-utilised plant which grows wild at roadside, on dry rocky slopes of stony soils and can withstand extreme temperatures ( $-30^{\circ}\text{C}$  to  $+35^{\circ}\text{C}$ ) of Ladakh and is also highly drought tolerant. This plant has multiple uses in cuisine as salad, pickle, and condiments<sup>5,6</sup>. *Capparis* is known to contain a wide variety of antioxidant compounds including

phenolic compounds which are found to be well correlated with antioxidant potential<sup>7</sup>.

Earlier phytochemical studies have shown the presence of alkaloids, flavonoids, polyphenols, indolic and aliphatic glucosinolates in *C. spinosa*<sup>8-10</sup>. It has also been studied pharmacologically, the alcoholic and aqueous extract of plant has antimicrobial, hepatoprotective, antihyperglycemic, and protects against oxidative stress in systemic sclerosis dermal fibroblasts<sup>11,12</sup>. The methanolic extract of leaves shows very high antioxidant activity and polyphenolic content<sup>7</sup>. The flower buds showed antioxidant and antibacterial activity and have excellent photoprotection against UVB-induced skin damage<sup>13</sup>. Natural antioxidants present in *C. spinosa* can scavenge harmful free radicals from our body. It is possible to reduce the risk of chronic diseases and prevent disease progression by either enhancing the body's natural antioxidant defense or by supplementing with proven dietary antioxidants<sup>14</sup>.

Until now, there has been no report on comparative study of the polyphenol contents and antioxidant activity in different edible parts of *C. spinosa* from trans-Himalayan region of Ladakh, India. Keeping in view to validate the medicinal potential of *C. spinosa* growing wild in trans-Himalayas, the present investigation was designed to study the major phytochemicals, polyphenols and antioxidant capacity among different edible parts viz. leaves, flower buds, fruits and roots of *C. spinosa* and to correlate the antioxidant capacity with that of polyphenol contents in different edible parts of *C. spinosa*.

## 2. MATERIALS AND METHODS

### 2.1 Plant Material and Preparation of Crude Extract

#### Extract

The edible parts used in the present investigation were collected from *C. spinosa* plants growing wild in Ladakh. Ladakh is a part of Indian trans Himalaya situated at an altitude of 2200 m - 6100 m amsl, is characterised by diverse and complex land formations. It is located at the latitude of 31°44'57" to 32°59'57"N and longitude of 76°46'29" to 78°41'34" E<sup>15</sup>. The plant was identified by its vernacular name by the local people and later authenticated by the Herbarium of Defence Institute of High Altitude Research, Leh-Ladakh, India. The edible parts were first decontaminated by washing under tap water followed by sterilised distilled water, shade dried at room temperature (26 °C) to constant weights. The dried samples (bulk of five samples) were then pulverised individually and 10 g each was separately shaken in methanol for 72 h in an orbital shaker at room temperature. Extracts were then filtered using a Buckner funnel and Whatman No. 1 filter paper and filtrate was concentrated to dryness under reduced pressures (337 mbar) at 40 °C using rotary evaporator. Then, the extract was resuspended in methanol to make 50 mg/ml stock solution<sup>16</sup>.

### 2.2 Chemicals

The chemicals of various brands used include 2,2-Diphenyl-2-picrylhydrazyl (DPPH), 2,2'-azinobis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS), 2,4,6-tripyridyl-s-triazine (TPTZ), potassium ferricyanide, potassium persulfate, trichloroacetic acid, gallic acid and quercetin from Sigma Chemical Co. (USA); FeCl<sub>3</sub>, ascorbic acid, and butylated hydroxyl toluene (BHT) from HIMEDIA Laboratories Pvt. Ltd. (India); Folin-Ciocalteu phenol reagent, anhydrous sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>), hydrochloric acid (HCl), glacial acetic acid, potassium chloride, sodium acetate trihydrate and solvent methanol were from Merck Chemicals (Darmstadt, Germany). All the chemicals used were of analytical grade.

### 2.3 Phytochemical Screening

Phytochemical analysis of the crude extract of *C. spinosa* was performed qualitatively for the presence of alkaloids, tannins, saponins and anthraquinones as plant constituents according to a standard method<sup>17</sup>. The alkaloids were detected by treating the samples with Dragendorff's reagent, which resulted in the formation of a precipitate at the base of the test tube. Those samples, in which green or black color appeared when mixed with aqueous FeCl<sub>3</sub>, were considered positive for tannins. Further, the presence of saponins was considered on the basis of frothing after vigorous shaking of diluted samples.

### 2.4 Determination of Antioxidant Activity

#### 2.4.1 DPPH Radical Scavenging Assay

The effect of the extracts on DPPH radical (DPPH<sup>+</sup>) was estimated using the standard method<sup>18</sup>. For this, the absorbance of the solution was measured spectrophotometrically at 517 nm using BHT as reference. The ability to scavenge DPPH<sup>+</sup> was calculated by the following equation: DPPH<sup>+</sup> scavenging activity (percent) =  $[(Abs_{Control} - Abs_{Sample}) / (Abs_{Control})] \times 100$

where Abs<sub>Control</sub> is the absorbance of DPPH<sup>+</sup> methanol; Abs<sub>Sample</sub> is the absorbance of DPPH<sup>+</sup> + sample extract /standard.

#### 2.4.2 ABTS Radical Scavenging Assay

ABTS radical (ABTS<sup>+</sup>) scavenging assay was determined as per Re<sup>19</sup>, *et al.* and the samples extract (1 ml) was mixed with 1 ml of the ABTS solution and the absorbance was taken at 734 nm after 7 min using spectrophotometer. The ABTS scavenging capacity of the extract was compared with that of BHT and per centage inhibition was calculated as ABTS radical scavenging activity (percent) =  $[(Abs_{Control} - Abs_{Sample}) / (Abs_{Control})] \times 100$  where Abs<sub>Control</sub> is the absorbance of ABTS radical+methanol; Abs<sub>Sample</sub> is the absorbance of ABTS radical+sample extract/standard. IC<sub>50</sub> (Inhibition coefficient) value was determined from the plotted graph of scavenging activity against concentration of all edible parts, which is defined as the amount of antioxidant necessary to decrease the initial DPPH/ABTS radical concentration by 50 per cent.

#### 2.4.3 Ferric Reducing Antioxidant Power Assay

The Ferric Reducing Antioxidant Power (FRAP) assay was conducted using method of Wong<sup>20</sup>, *et al.* The increase in absorbance was measured using spectrophotometer at 593 nm. The per cent of antioxidant was calculated using the formula, per cent of antioxidant (percent) =  $[(Abs_{Sample} - Abs_{Control}) / Abs_{Sample}] \times 100$ .

#### 2.4.4 Determination of Total Phenols

Total phenol contents in the extracts were determined by using modified Folin-Ciocalteu method<sup>21</sup>. The absorbance of the solution was measured at 765 nm. Total phenolic content was expressed as mg/g tannic acid equivalent using the following equation based on the calibration curve:  $y = 0.1216x$  (R<sup>2</sup>=0.9365), where x was the absorbance and y was the gallic acid equivalent (mg/g).

#### 2.4.5 Estimation of Total Flavonoids

Estimation of the total flavonoids in the caper extracts was carried out as per Ordon<sup>22</sup>, *et al.* and absorbance was measured at 420 nm. The yellow color indicated the presence of flavonoids and total flavonoid was calculated as quercetin equivalent (mg/g) using the following equation based on the calibration curve:  $y = 0.0255x$  (R<sup>2</sup>=0.9812), where x was the absorbance and y was the quercetin equivalent (mg/g).

## 2.5 Statistical Analysis

The experimental results were expressed as mean ± standard deviation (SD) of three replicates ( $n = 3$ ) and the data was subjected to one-way analysis of variance (ANOVA) using SPSS 11.5. Duncan's multiple range tests were used to assess difference between means. Regression test was used to assess correlation between means. The dendrogram was made using Ward method and distance is expressed as Euclid distance. P-values < 0.05 were regarded as significant.

## 3. RESULT AND DISCUSSION

### 3.1 Phytochemical Screening

The phytochemical screening of *C. spinosa* extracts

showed the presence of different chemical classes, such as alkaloids and saponins. The literature has also shown the presence of various plants constituents and the presence of these phytochemicals in *C. spinosa* may be responsible for its usefulness in various diseases. Though, further studies are required to know the precise nature of chemical constituents mediating alleged biological activities.

### 3.2 DPPH Radical Scavenging Activity

The DPPH radical scavenging activity provides information on their activity of the test compounds with a stable free radical. DPPH is one of the compounds that possess a proton free radical and shows absorption band at 517 nm in visible region. Methanolic extracts of leaves showed the highest scavenging effect (70.86 per cent) at a concentration of 0.1 mg/ml, whereas methanolic extract of dried fruits exhibited the lowest activity at the same concentration. Among all edible parts, though the DPPH radical scavenging abilities of the extracts were less than BHT (72.09 per cent).

IC<sub>50</sub> value was deciphered from the graph of scavenging ability against the concentration of methanolic extract of *C. spinosa*. Higher IC<sub>50</sub> value indicated lower antioxidant activity and *vice-versa*. Table 1 showed the highest IC<sub>50</sub> value for DPPH in dried fruits (0.097 mg/ml) and lowest in leaf sample (0.050 mg/ml). Further, the degree of reduction in absorbance indicates the radical scavenging power of the extract. The effect of antioxidants on DPPH is probably due to its hydrogen donating ability<sup>23</sup>. Though the DPPH radical scavenging abilities of the extracts were less than BHT, the study showed that the extracts have the proton-donating ability and could serve as free radical inhibitors or scavengers, acting possibly as primary antioxidants. This study suggests that all edible parts of *Capparis* plant possess antioxidant activity (Fig. 1).

### 3.3 ABTS Radical Scavenging Activity


ABTS, a protonated radical, has characteristic absorbance maxima at 734 nm which decreases with the scavenging of the proton radicals<sup>24</sup>. The methanol extracts of the leaves of *C. spinosa* were fast and effective scavengers of the ABTS radical and this activity was comparable to that of ascorbic acid and BHT. At concentration of 0.02 mg/ml, 0.04 mg/ml, 0.06 mg/ml, and 0.08 mg/ml the ascorbic acid and BHT exhibited higher activity than the leaves extracts, but at 0.1 mg/ml the activity of the leaves extracts were similar to that of ascorbic acid and BHT (100 per cent). Lowest activity found in dried fruit extract (61.15 per cent) at a concentration of 0.1 mg/ml.

Table 1 revealed that highest IC<sub>50</sub> value for ABTS found in dried fruits (0.086 mg/ml) and lowest were found in leaves sample (0.033 mg/ml). The scavenging of the ABTS<sup>+</sup> by the extracts was found more than that of DPPH radical. Factors like stereoselectivity of the radicals or the solubility of the extract in different testing systems have been reported to affect the capacity of extracts to react and quench different radicals<sup>25</sup>. Some compounds which have ABTS<sup>+</sup> scavenging activity did not show DPPH scavenging activity<sup>26</sup>, but same was not found true for this study (Fig. 1).

**Table 1.** Free radical scavenging activity (IC<sub>50</sub>) value for methanolic extract of all edible parts of *C. spinosa* collected from Ladakh region. [Values are expressed as mean ± SD, n = 3]

Edible parts	IC <sub>50</sub> mg/ml	
	DPPH	ABTS
Leaves	0.050 ± 0.003 <sup>a</sup>	0.033 ± 0.003 <sup>a</sup>
Flower Buds (6-8 mm)	0.068 ± 0.002 <sup>b</sup>	0.048 ± 0.002 <sup>b</sup>
Flower Buds (9-12 mm)	0.091 ± 0.002 <sup>c</sup>	0.077 ± 0.002 <sup>d</sup>
Roots	0.094 ± 0.003 <sup>cd</sup>	0.066 ± 0.003 <sup>c</sup>
Fruits (Dried)	0.097 ± 0.002 <sup>d</sup>	0.086 ± 0.002 <sup>c</sup>

Values with different superscripts in a column differ significantly ( $p < 0.05$ )



**Figure 1.** Free radicals scavenging activity determined with ABTS, DPPH and FRAP assays at 0.04 mg/ml concentration of methanolic extract of edible parts of *C. spinosa* samples.

### 3.4 FRAP

FRAP assay measures the reducing potential of an antioxidant reacting with a ferric tripyridyltriazine (Fe<sup>3+</sup>-TPTZ) complex and producing a colored ferrous tripyridyltriazine (Fe<sup>2+</sup>-TPTZ)<sup>27</sup>. (Fig. 2) showed that methanolic extract of *Capparis* leaves had highest total antioxidant potential (73.54 percent to 87.14 percent) compared to BHT (67.51 percent to 86.97 percent). As per the FRAP assay, the minimum antioxidant activity was recorded in dried fruit extract (64.70 percent to 81.81 percent), however, other edible parts are not significantly different in its antioxidant contents. Generally, the reducing properties of any sample are associated with the presence of those compounds, which can break the free radical chain by donating a hydrogen atom<sup>28</sup>. In this study, phenolic compounds of all edible parts of *Capparis* exhibited high reducing power on Fe<sup>3+</sup>-TPTZ (Fig 1). Due to its redox properties, phenolic compounds act as reducing agents, hydrogen donors and singlet oxygen quenchers. The redox potential of phenolic compounds plays an important role in determining the antioxidant capacity<sup>29</sup>.

### 3.5 Total Phenols

Polyphenolic compounds are known to have antioxidant activity<sup>30</sup>. This activity is believed to be mainly due to their redox properties, which play an important role in adsorbing and neutralizing free radicals, quenching singlet and triplet oxygen, or decomposing peroxides<sup>31</sup>. The content of phenol compounds determined by the Folin-Ciocalteu method for the *C. spinosa* leaves analysed is shown in (Fig. 3). Total phenolic

compounds ranged from 4.07 to 24.78 mg GAE/g dry wt. The highest total phenolic content was found in the leaves (24.78) whereas the lowest phenolic content was found in the dried fruits (4.7 mg GAE/g dry wt).




Figure 2. Antioxidant content (%) of methanolic extract of all edible parts of *Capparis spinosa* expressed as per cent of antioxidant using FRAP method.

In fact, many medicinal plants contain large amounts of antioxidants and many of these phytochemicals possess significant antioxidant capacities that are associated with lower occurrence and lower mortality rates of several human diseases<sup>32</sup>.

### 3.6 Total Flavonoids

Fig. 3 exhibits the flavonoid contents of all edible parts of *C. spinosa*. The result revealed that leaves having maximum flavonoids contents (5.69 mg quercetin/g dry wt.) and dried fruits having no flavonoid contents among all edible parts. This result strongly suggests that polyphenols are important components of *Capparis*, which are responsible for not only its antioxidant activities but some of its pharmacological effects, could be attributed to the presence of these valuable constituents. The presence of flavonoids might also influence the antioxidant capacity<sup>33</sup>.




Figure 3. Total phenolic content (gray bars, mg of GAE g<sup>-1</sup> of DW) and total flavonoid content (white bars, mg of quercetin g<sup>-1</sup> of DW) of *C. spinosa* edible parts.

### 3.7 Correlation

Several studies have reported correlations between the antioxidant activities measured by different methods, as well as the correlations between those methods and phytochemical concentrations in various food commodities<sup>34</sup>. However, this type of information is very limited for *C. spinosa*. The IC<sub>50</sub> value for DPPH was highly correlated with that of ABTS

(R<sup>2</sup>=0.9084) and FRAP assay (R<sup>2</sup> = 0.9771). The result suggested that the three methods have similar predictive capacity for antioxidant activities of *C. spinosa*. However, IC<sub>50</sub> value of ABTS is reasonably correlated with FRAP assay (R<sup>2</sup> = 0.5838). There was a high correlation (R<sup>2</sup> = 0.90) between ABTS and DPPH values for various fruit extracts<sup>35</sup>.

Several studies have shown correlation between antioxidant activity and total phenolic contents<sup>7</sup>. The total phenolic contents were highly correlated with IC<sub>50</sub> value of ABTS (R<sup>2</sup>= 0.9084), DPPH (R<sup>2</sup>= 0.9388) and FRAP value (R<sup>2</sup> = 0.9618). However, total flavonoid contents were reasonably correlated with ABTS (R<sup>2</sup>= 0.7449) and DPPH (R<sup>2</sup>= 0.8791) but highly correlated with FRAP value (R<sup>2</sup> = 0.9588).

### 3.8 Cluster Analysis

Cluster analysis of *C. spinosa* edible parts collected from Ladakh region showed that cluster based on total antioxidant activity (DPPH, ABTS, and FRAP) was almost similar to cluster based on total phenolic contents (Figs. 4 and 5)<sup>36</sup>. It revealed that in case of capers, total Antioxidant capacity is equally contributed by total phenolics and total flavonoids. Antioxidant activities varied widely among all the edible parts of *C. spinosa*. There were good correlations between the




Figure 4. Dendrogram of different edible parts of *C. spinosa* according to cluster analysis of similarity on the basis of total AOA determined by DPPH, ABTS and FRAP assays using Ward method.




Figure 5. Dendrogram of different edible parts of *C. spinosa* according to cluster analysis of similarity on the basis of total phenolic and flavonoid contents using Ward method.

antioxidant activities measured by DPPH, ABTS, and FRAP as well as total phenolic contents, suggesting that these methods have similar predictive capacity for antioxidant activities of edible parts of *C. spinosa*. High correlation between the DPPH, ABTS, FRAP and phenol as well as flavonoid contents indicated that the total phenolic contents can be used as indicator for antioxidant activities of edible parts of *C. spinosa*. This result again suggests that Antioxidant capacity of capers is caused mainly by phenolics as well as flavonoids.

#### 4. CONCLUSION

The results revealed that the methanolic extracts of *C. spinosa* leaves possess a strong antioxidant/free radical scavenging activities among all the edible parts, which is probably due to the presence of high concentration of polyphenolic compounds. The strong antioxidant activity of all edible parts of *C. spinosa* may, therefore, be a good candidate for functional foods as well as plant-based pharmaceutical products. Further studies are required to identify the active principle responsible for the significant antioxidant effect.

#### REFERENCES

- Katalinic, V.; Milos, M.; Modun, D.; Music, I. & Boban, M. Antioxidant effectiveness of selected wines in comparison with (+) catechin. *Food Chem.*, 2004, **86**, 593–600.  
doi: 10.1016/j.foodchem.2003.10.007
- Mishra, G.P.; Singh, N.; Kumar, H. & Singh, S.B. Protected cultivation for food and nutritional security at Ladakh. *Def. Sci. J.*, 2010, **61**(2), 219-225.
- Mishra, G.P.; Singh, R.; Bhojar, M.S. & Singh, S.B. *Capparis spinosa*: Unconventional potential food source in cold arid deserts of Ladakh. *Curr. Sci.*, 2009, **96**, 1563–1564.
- Bhojar, M.S.; Singh, R.; Mishra, G.P. & Stobdan, T. Lesser-known plants to enrich vegetable basket in Ladakh. *Indian Hort.*, 2009, **54**(4), 16-17.
- Bhojar, M.S.; Mishra, G.P.; Singh, R. & Singh, S.B. Effect of various dormancy breaking treatments on the germination of wild caper (*Capparis spinosa* L.) seeds from the cold arid desert of trans-Himalayas. *Indian J. Agr. Sci.*, 2010, **80**(7), 620-624.
- Bhojar, M.S.; Mishra, G.P.; Naik, P.K.; Murkute, A.A. & Srivastava, R.B. Genetic variability studies among natural populations of *Capparis spinosa* from cold arid desert of trans-Himalayas using DNA markers. *Natl. Acad. Sci. Lett.*, 2012, **35**(6), 505-515.  
doi: 10.1007/s40009-012-0086-y
- Bhojar, M.S.; Mishra, G.P.; Naik, P.K. & Srivastava, R.B. Estimation of antioxidant activity and total phenolics among natural populations of *Capparis spinosa* leaves collected from cold arid desert of trans-Himalayas. *Aust. J. Crop Sci.*, 2011, **5**(7), 912-919.
- Sharaf, M.; El-Ansari, M.A. & Saleh, N.A. Quercetin triglycoside from *Capparis spinosa*. *Fitoterapia*, 2000, **71**, 46-49.  
doi: 10.1016/S0367-326X(99)00116-1
- Ulukapı, K.; Buse O.; Kulcan, A.A.; Tetik, N.; Ertekin, C. & Onus, A.N. Evaluation of biochemical and dimensional properties of naturally grown *Capparis spinosa* var. *spinosa* and *Capparis ovata* var. *palaestina*. *Int. J. Agril. Inn. Res.*, 2016, **5**(2), 195-200.
- Tagnaout, I.; Zerkani, H.; Mahjoubi, M.; Bourakhouadar, M.; Alistiqsa, F.; Bouzoubaa, A. & Zair, T. Phytochemical study, antibacterial and antioxidant activities of extracts of *Capparis spinosa* L. *Int. J. Pharmacol. Phytochem. Res.*, 2016, **8**(12), 1993-2006.
- Feng, X.; Lu, J.; Xin, H.; Zhang, L.; Wang, Y. & Tang, K. Anti-arthritis active fraction of *Capparis spinosa* L. fruits and its chemical constituents. *Yakugaku Zasshi*, 2011, **131**(3), 423-429.  
doi: 10.1248/yakushi.131.423
- Anwar, F.; Muhammad, G.; Hussain, M.A.; Zengin, G.; Alkharfy, K.M.; Ashraf, M. & Gilani, A.H. *Capparis spinosa* L.: A plant with high potential for development of functional foods and nutraceuticals/ pharmaceuticals. *Int. J. Pharmacol.*, 2016, **12**, 201-219.  
doi: 10.3923/ijp.2016.201.219
- Bouriche, H.; Karnouf, N.; Belhadj, H.; Dahamna, S.; Harzalah, D. & Senator, A. Free radical, metal-chelating and antibacterial activities of methanolic extract of *Capparis spinosa* buds. *Adv. Environ. Biol.*, 2011, **5**(2), 281-287.
- Stanner, S.A.; Hughes, J.; Kelly, C.N. & Buttriss, J.A. Review of the epidemiological evidence for the antioxidant hypothesis. *Public Health Nutrition*, 2000, **7**, 401-422.  
doi: 10.1079/PHN2003543
- Bhojar, M.S.; Mishra, G.P.; Singh, R. & Singh, S.B. Ethno-botany of traditional wild edible plants from cold arid desert of Ladakh-potential source of winter vegetables. *The Indian Forester*, 2011, **137**(8), 1029-1033.
- Navaie, B.A.; Kavosian, S.; Fattahi, S.; Hajian-Tilaki, K.; Asouri, M.; Bishekolaie, R. & Akhavan-Niaki, H. Antioxidant and cytotoxic effect of aqueous and hydroalcoholic extracts of the *Achillea millefolium* L. on MCF-7 breast cancer cell line. *Int. Biol. Biomed. J.*, 2015, **1**(3), 119-125.
- Evans, W.C. Phytochemistry. In Trease and Evans Pharmacognosy, 5th Edn, Elsevier: Delhi, 2006, 135–150.
- Liyana-Pathiranan, C.M. & Shahidi, F. Antioxidant activity of commercial soft and hard wheat (*Triticum aestivum* L) as affected by gastric pH conditions. *J. Agric. Food Chem.*, 2005, **53**, 2433-2440.  
doi: 10.1021/jf049320i
- Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M. & Rice-Evans, C. Antioxidant activity applying an improved ABTS radical cation decolorisation assay. *Free Radical Bio. Med.*, 1999, **26**, 1231-1237.  
doi: 10.1016/S0891-5849(98)00315-3
- Wong, S.P.; Lai, P.L. & Jen, H.W.K. Antioxidant activities of aqueous extracts of selected plants. *Food Chem.*, 2006, **99**, 775-783.

- doi: 10.1016/j.foodchem.2005.07.058
21. Wolfe, K.; Wu, X. & Liu, R.H. Antioxidant activity of apple peels. *J. Agric. Food Chem.*, 2003, **51**, 609-614.  
doi: 10.1021/jf020782a
  22. Ordon, E.A.; Gomez, J.D.; Vattuone, M.A. & Isla, M.I. Antioxidant activities of *Sechium edule* (Jacq.) Swart extracts. *Food Chem.*, 2006, **97**, 452-458.  
doi: 10.1016/j.foodchem.2005.05.024
  23. Baumann, J.; Wurn, G. & Bruchlausen, F.V. Prostaglandin synthetase inhibiting O-2 radical scavenging properties of some flavonoids and related phenolic compounds. In Deutsche Pharmakologische Gesellschaft Abstracts of the 20th spring meeting, Naunyn-Schmiedebergs Abstract No: R27 cited in Arch. Pharmacol., 1979, **307**, R1 –R77.
  24. Mathew, S. & Abraham, E.T. In vitro antioxidant activity and scavenging effects of *Cinnamomum verum* leaf extract assayed by different methodologies. *Food Chem. Toxicol.*, 2006, **44**, 198-206.  
doi: 10.1016/j.fct.2005.06.013
  25. Yu, L.; Haley, S.; Perret, J.; Harris, M.; Wilson, J. & Qian, M. Free radical scavenging properties of wheat extracts. *J. Agric. Food Chem.*, 2002, **50**, 1619–1624.  
doi: 10.1021/jf010964p
  26. Wang, M.; Li, J.; Rangarajan, M.; Shao, Y.; La, V.E.J.; Huang, T. & Ho, C. Antioxidative phenolic compounds from Sage (*Salvia officinalis*). *J. Agric. Food Chem.*, 1998, **46**, 4869–4873.  
doi: 10.1021/jf980614b
  27. Benzie, I.F. & Strain, J.J. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Method Enzymol.*, 1999, **299**, 15–27.  
doi: 10.1016/S0076-6879(99)99005-5
  28. Duh, P.D.; Du, P.C. & Yen, G.C. Action of methanolic extract of mung hulls as inhibitors of lipid peroxidation and non-lipid oxidative damage. *Food Chem. Toxicol.*, 1999, **37**, 1055–1061.  
doi: 10.1016/S0278-6915(99)00096-4
  29. Rice-Evans, C.A.; Miller, N.T. & Paganga, G. Antioxidant properties of phenolic compounds. *Trends Plant Sci.*, 1997, **4**, 304–309.  
doi: 10.1016/S1360-1385(97)01018-2
  30. Tepe, B.; Sokmen, M.; Akpulat, H.A. & Sokmen, A. Screening of the antioxidant potentials of six *Salvia* species from Turkey. *Food Chem.*, 2006, **95**, 200-204.  
doi: 10.1016/j.foodchem.2004.12.031
  31. Zheng, W. & Wang, S.Y. Antioxidant activity and phenolic compounds in selected herbs. *J. Agric. Food Chem.*, 2001, **49**, 5165–5170.  
doi: 10.1021/jf010697n
  32. Djeridane, A.; Yousfi, M.; Nadjemi, B.; Boutassouma, D.; Stocker, P. & Vidal, N. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem.*, 2006, **9**, 654-660.  
doi: 10.1016/j.foodchem.2005.04.028
  33. Kaur, C. & Kapoor, H.C. Review: Antioxidants in fruits and vegetables – the millennium’s health. *Int. J. Food Sci. Technol.*, 2001, **36**, 703–725.  
doi: 10.1111/j.1365-2621.2001.00513.x
  34. Awika, J.M.; Rooney, L.W.; Wu, X.; Prior, R.L. & Cisneros-Zevallos L. Screening methods to measure antioxidant activity of sorghum (*Sorghum bicolor*) and sorghum products. *J. Agric. Food Chem.*, 2003, **51**, 6657–666.  
doi: 10.1021/jf034790i
  35. Leong, L.P. & Shui, G. An investigation of antioxidant capacity of fruits in Singapore markets. *Food Chem.*, 2001, **76**, 69–75.  
doi: 10.1016/S0308-8146(01)00251-5
  36. Kumar, J.; Bajaj, P.; Mishra, G.P.; Singh, S.B.; Singh, S.B. & Naik, P.K. Utilisation of EST-derived SSRs in the genetic characterisation of *Artemisia annua* L. genotypes from Ladakh, India. *Indian J. Biotechnol.*, 2014, **13**, 464-472.

#### ACKNOWLEDGMENT

Authors thank Director, DIHAR, Leh for providing necessary facilities for this study. This study was funded by the Defence Research and Development Organisation, Ministry of Defence, India.

#### CONTRIBUTORS

**Dr Manish S. Bhoyar** has received his PhD from Jaypee University of Information Technology, Solan, in 2012 and his major research area is genetic improvement of vegetables. Currently working as Fellow Scientist at National Innovation Foundation-India, Gandhinagar, Gujarat.

He was involved in the execution of the laboratory experiments and data analysis.

**Dr Gyan P. Mishra** has received his PhD from, Indian Agricultural Research Institute, New Delhi, in 2006 and his major research area is genetic improvement and molecular breeding of crop plants. Currently working as Senior Scientist (Genetics and Plant Breeding) at Indian Agricultural Research Institute, New Delhi.

He was involved in the conceptualisation of experiment, data interpretation and manuscript writing.

**Dr Pradeep K. Naik** has received his PhD from Sambalpur University, Odisha, in 2001 and his major research area is genetic improvement of high altitude pants. Currently he is working as Professor at Sambalpur University, Odisha.

He was involved in the data analysis and manuscript editing.

**Dr Shashi Bala Singh** has received her PhD from All India Institute of Medical Sciences, New Delhi, in 1986. She has more than 100 publications in journals. Currently she is working as Director General-Life Sciences (LS), DRDO HQrs.

She was involved in the conceptualisation of experiment and data interpretation.