

REVIEW PAPER

Nutritional Intervention during Chemical, Biological, Radiological, Nuclear Environments: A Dietary Perspective

Dev Kumar Yadav, Janifer Raj Xavier*, Om Prakash Chauhan,
Prakash Eknath Patki, and Rakesh Kumar Sharma

DRDO-Defence Food Research Laboratory, Mysore, India-570011.

*E-mail: janifer@dfrl.drdo.in

ABSTRACT

The future war scenario is based on use and applications of various conventional and non-conventional agents which includes weaponised or non-weaponised chemical, biological, radiological, nuclear (CBRN), toxic industrial materials, direct energy devices/ weapons, and or high yield explosives. These include nerve agents, blood agents, vesicants or skin blistering agents, lung irritants, asphyxiants or choking agents. Biological weapons are basically disease causing microorganisms and other replicating entities including viruses, infectious nucleic acids and prions. These agents have ability to infect host and are highly virulent, pathogenic and dangerous in nature. The interface between ammunitions and above agents is soldier whose physical and mental health is affected as enough precautionary measures are not adopted. The reducing environment thus created has various agents which enter into exposed body and lead to mild to serious damage to various vital parts of the human body. As food is important component for survival and intrinsic to basic human nutrition and health, therefore, it is imperative to develop certain kind of a wholesome meal system which can be consumed by the soldiers tasked with combating CBRN situations during such operations. Such meals can be in the form of solid or liquid type and packaged in suitable delivery system, compatible and amenable with the CBRN suit. Food can be contaminated during CBRN conditions by coming in direct or indirect contact with CBRN agents. Therefore, the food materials to be used under such conditions need to be protected in suitable coverings as consumption of contaminated food can be lethal. Designer meal for CBRN environment is not only suitable for soldiers but also to all human interface dealing with similar scenario *viz.* the low intensity conflicts and surgical operations, nuclear submarines, cosmonauts, pilots, individuals handling radiation equipment and patient undergoing chemotherapy for cancer.

Keywords: Chemical; Radiological; Biological; CBRN meal; Protection

1. INTRODUCTION

The technological advancement in science has opened entirely new avenues for every spheres of life *viz.* information technology, pharmaceutical, food, defence etc. The national security for any nation is of prime concern and this is why the forthcoming war patterns indicate the use of both conventional and unconventional weapons which are deadly threat to the mankind. It may be in form of a chemical, biological, radiological and or nuclear (CBRN) entity which commits to severely affect the whole humanity^{1,2}. In such scenario, the fighting soldiers and the exposed population must have proper and enough defence measures for safety of individual inclusive of physical, biological, and psychological protection¹⁻³. Combat feeding of troops under varied climatic, terrain and weapon platform conditions is the highlight and mainstay of R&D programmes of Defence Research and Development Organisation (DRDO) in general and Defence Food Research Laboratory (DFRL) in particular. Damages to biological system by ionising radiation are either by directly affecting bio-macromolecules or by generation of highly reactive free radicals. Free radicals cause

oxidative damages to bio molecules like lipid, protein and DNA such as single strand double strand breaks and DNA-DNA and DNA-Protein cross linkage etc. Rate of damage also increases in a dose dependent manner, repair mechanism of cell will try to resolve situation, but most of them will be result in unrepaired or mis repaired stage that will lead to mutation or cell death⁴. Many studies are going on for the development of radio protectors to save our biological system from the harmful effects of radiation. That resulted in development of a number of compounds of synthetic origin such as vitamins, glycosides, nucleic acid derivatives and phosphorothioates such as amifostine. Radio protectors action can be summarised as suppression of free radicals, detoxifying reactive species and enhancing action of superoxide dismutase prostaglandins interleukins etc. stimulating DNA repair and suppressing cell division⁵.

But still problems persist in form of fewer efficacies and associated toxicity with their effective doses. As an alternative source many plants have been studied and proved for their radio protective efficiency. Studies are available for the radio protective properties of plants such as *Podophyllum hexandrum*, *Spirulina plantensis*, *Chlorella vulgaris*, *Centella asiatica*,

Tinosora cordifolia etc. Plants with active constituents such as silibinin, caffeic acid, thymol, phenylpropanoids phenyl esters, phycocyanin, chlorogenic acid, curcumin lycopene, gingerol, quercetin, resveratrol, orientin, vicenin, vitamin C and vitamin E, are reportedly shown radioprotective properties^{6,7}. Major advantage of natural compounds are their less toxicity when they are administered before or after irradiation shows protective actions⁸. Several plant based compounds are already using widely for treatment of cancer in clinical field e.g. Taxol, vinblastine, vinca alkaloids, topotecan and vincristine⁹. This review elucidates a general understanding of nutritional interventions and their role in reducing the ill effects of radiation on the soldier exposed to CBRN scenario.

2. CBRN ENVIRONMENT

An apex body entitled ONTARIO explains CBRN as weaponised or non-weaponised chemical, biological, radiological and nuclear materials that can cause great harm and pose significant threats in the hands of terrorists. Weaponised materials can be delivered using conventional bombs (e.g., pipe bombs), improved explosive materials (e.g., fuel oil-fertiliser mixture), and enhanced blast weapons (e.g., dirty bombs)^{2,3}. Non-weaponised materials are traditionally referred to as dangerous goods (DG) or hazardous materials (HAZMAT) and can include contaminated food, livestock, and crops. An accidental CBRN incident is an event caused by human error or natural or technological reasons, such as spills, accidental releases, or leakages. These accidental incidents are usually referred to as DG or HAZMAT accidents. Outbreaks of infectious diseases, such as SARS, or pandemic influenza are examples of naturally occurring biological incidents¹⁰⁻¹¹.

3. EXPOSURE AND HEALTH IMPLICATIONS

There are multiple harmful effects of radiation, and many of them go unnoticed in the body. A low level exposure can lead to major digestive imbalance, blood alteration and even the destruction of many cellular structures in the body's key organs and tissue systems. Common signs of low level radiation exposure include symptoms such as fatigue, headaches, nausea, scalp tenderness, scalp discoloration, and dry/itchy skin^{12,13}. In extreme cases, low exposure of radiation may also cause brain damage, memory problems, mood changes, and reduced listening capacities, psychomotor abilities, and information processing times. Large amounts of radiation exposure can have devastating consequences such as bloody vomit, hair loss, nerve damage, blood vessel destruction, seizures and, with prolonged strong exposure, can even lead to death¹³⁻¹⁶. Our circulatory system is also vulnerable to radiation. Even very low level of radiation exposure reduces lymphocytes in the blood. When lymphocytes are lowered, we are more likely to get infections, and what is known as mild radiation sickness. Symptoms are flu-like and may increase the likelihood of developing leukemia and lymphoma in future. The reproductive system is also vulnerable to exposure. Long-term radiation exposure can even lead to sterility, birth defects, and still-births¹³⁻¹⁷.

Ionising radiations either directly affects the bio-macromolecules and cause damage or it will generate free radicals by reacting with smaller molecules like water.

Reaction proceeds in a dose dependent manner and self defence mechanism of cells fail when dose goes high. Reactive oxygen species generated as a result of radiolysis includes hydrated electron (e_{aq}^-), hydrogen radical ($\cdot H$), hydroxyl radical ($\cdot OH$), peroxy radical ($\cdot ROO$), superoxides ($O_2^{\cdot -}$)¹⁸. Free radicals react with cellular structures like DNA, lipid membranes, and proteins cause's damage. Self defence mechanism of cells will try to mitigate these free radicals by using enzymes such as Glutathione peroxidase reduces hydroxyl radicals and generates water, superoxide dismutase converts superoxide radicals into hydrogen peroxide and later catalase reduces them to water. But when these radicals are produced at higher rate cells will not be able to control their production and get injured. Direct effects will result in single and double strand breaks, cell try to repair these damages by a series of enzymatic reactions; either it will result in successful repair or to induce apoptosis¹⁹. Suppression and damages to the haematopoietic system is one of the most dangerous consequences of radiation exposure. Ionising radiation can affect directly or indirectly haematopoiesis at various levels such as stem cells, blood cells or progenitor cells etc. Directly it will cause damage to stem cells or progenitor cells and indirectly it can interfere with the communication pathways between the cells and the microenvironment. Radioprotective compounds from herbal sources can stimulate generation of haematopoietic elements thereby reduce the harmful effects of radiation. Thus, nuclear accidents results in a massive destruction to the humanity. When exposed to irradiation all the systems controlling growth and maintenance of the cell get damaged so that cell will not be able to function normally. So that an ideal radio protector should improve wound healing activity of the cells so that they can recover fast. Mechanism of wound healing includes mainly three phase namely inflammation, proliferation and remodelling.

4. DIETARY SUPPLEMENTS AND RATIONALE

Certain foods play crucial role in protecting the exposed body against radiation and these facts came out of the various studies carried out after Chernobyl disaster as shown in Table 1. These are plant foods *viz.* fruits and vegetables which have cancer fighting active principal compounds. In a study, the Swedish government has measured the residual radiation level in food following the Chernobyl disaster and mentioned that animal foods such as meat, fish, and dairy products have higher amount of radioactive substances than the plant foods such as cereals, pulses, fruits, and vegetables as they poorly absorb radiation²⁰. The animal tissues are more receptive of radioactive doses than the plant matrix²¹.

The natural presence of phytochemicals inclusive of fibers, antioxidants, and pigments in foods provide them the antioxidant activity necessary to fight against cellular damage caused by free radicals formed during radiation exposure. Dietary fibers from lemon peel and beetroot do possess radioprotective properties and these fibers in concentrated form may be used to eliminate the nuclides and radioactive substances. Pectin naturally present in many fruits do provide protection from radiation in particular from Cs-137. A study evaluating

Table 1. Dietary supplements for protection from radiation exposure

Group	Nominations
Salts	Potassium Iodide It saturates the thyroid gland with iodine by competing with radioactive iodine ⁴⁴ .
Phytochemicals	Resveratrol, Quercetin, Epigallocatechin gallate, Isoflavon (genestein), Curcumin, N-acetyl cysteine (NAC-Sulphur containing compounds), Coenzyme Q ₁₀ , lipoic acid ⁴³ .
Herbs and Algae	<i>Ginkgo biloba</i> , Ginseng, <i>Spirulina</i> , <i>Chlorella</i> ⁴⁷ .
Enzymes	Superoxide dismutase (SOD), Catalase, Bowman-Birk inhibitor ⁴⁵ .
Micro elements	ACE Vitamins, Selenium, Zn, and Mg ⁴⁶ .

30,000 atomic bomb survivors in Hiroshima and Nagasaki found that the population heavily fed with fruits and vegetables had 13 per cent less risk of death from cancer and similar diseases. Sulfur containing antioxidants present in cruciferous vegetables such as broccoli, kale, and cabbage do protect from radiation exposure as they have high detoxifying properties²².

The foods with high antioxidant activity and water soluble vitamins are highly effective in combating the stress and the damage caused due to exposure in CBRN environment²³. Certain studies showed that naturally colorful fruits and vegetables juices boost non-specific immunity and helps in rejuvenating the body from the detrimental effect of CBRN environment. Prolonged exposure and high dose facing results into mutagenic effects of the radiation leads to benign and malignant cancers in the exposed individuals, many green leafy vegetables have active phytochemical principals that can cope up with these after affects. *Murraya koenigii* (curry) leaves was evaluated for its broad-spectrum antioxidant and anti-mutagenic activities and found quite effective. Antioxidant, anticarcinogenic, and anti-inflammatory properties of pomegranate constituents have been well studied and it established the fact that these are very effective in the treatment and prevention of cancer, cardiovascular disease, diabetes, dental conditions, erectile dysfunction, bacterial infections and antibiotic resistance, and ultraviolet radiation-induced skin damage. Phytochemical analysis by HPLC, LC-MS and total phenolic content revealed high content of ellagitannins which might be responsible for promising antioxidant and anti-mutagenic activities of *P. granatum* peel extract.

Antioxidant and anti-mutagenic potential of Guava (*Psidium guajava*) leaf extracts was also explored in previous studies. Guava contains high amount of phenolics responsible for broad-spectrum anti-mutagenic and antioxidant properties *in vitro* and could be potential candidates to be explored as modern phyto-medicine. Consumption of carotenoid-rich (lutein, zeaxanthin, beta-carotene) foods even for a short period of time gives protection against oxidative stress. The results obtained from the study suggest that the protective role is not specifically related to carotenoids. However, they may contribute together with other substances present in plant foods such as spinach, tomato etc., to lymphocyte resistance to oxidative damage. Lycopene acts synergistically, as an effective antioxidant against LDL oxidation, with several natural antioxidants such as vitamin E, the flavonoid glabridin, the phenolics rosmarinic acid and carnosic acid, and garlic²⁴. Microalgae are a potential novel source of bioactive molecules,

including a wide range of different carotenoids that can be used as nutraceuticals, food supplements, and novel food products. The potential of carotenoids from marine microalgae can be used as therapeutics to treat or prevent oxidative stress-related diseases^{25,26}. It is imperative to understand that the ascorbic acid, tocopherol, β carotene etc., have protective role against γ -radiation induced DNA damage in human lymphocytes. The radio-protective action of antioxidant Vitamins C, E, and beta-carotene were dependent upon their concentration as well as time and sequence of application²⁷ as shown in Table 1.

5. MODUS OPERANDI OF ANTI-INFLAMMATORY AND ANTIOXIDANT AGENTS IN TARGETING RADIATION INDUCED AGENTS

Cyclooxygenase-2 (COX-2) is an enzyme induced by a variety of factors such as tumor promoters, cytokines, growth factors and hypoxia. It is involved in metabolic conversion of arachidonic acid into endoperoxides which are used to synthesise prostaglandins, prostacyclin, or thromboxanes. The radiation exposure leads to excessive production of COX-mediated eicosanoids such as prostaglandins; thromboxane and 5-lipoxygenase (5-LOX) mediated leukotrienes. Eicosanoids serve as cell signalling molecules that are significant mediators of immune responses. The increased levels of prostaglandins and thromboxane, and thus an inflammatory state in the tissue, can arise within hours and persist up to weeks after irradiation in a wide range of organs and tissues. Traditional non-steroidal anti-inflammatory drugs (NSAIDs) exhibit analgesic and antipyretic effects by blocking COX-derived prostaglandin formation, with less adverse effects involving the gastrointestinal tract and reproductive system. Prostaglandins production in inflamed tissue down-regulates inflammation-related pathological symptoms such as pain and swelling on administration of COX-2 specific inhibitors. Natural antioxidants such as curcumin²⁸, asymmetrical indole curcumin analogs such as *N*-(3-(3-*1H*-indole-3-yl)acryloyl)phenyl)-4-fluorobenzamide, pyrazole and isoxazole²⁹, resveratrol, *trans*-3,5,4'-trihydroxystilbene³⁰, flavocoxid, an extract containing a mixture of catechin and baicalin, cardamonin³¹, flavones hesperidin and its aglycon hesperetin³², isorhamnetin, an antioxidant 3'-*O*-methylated metabolite of quercetin and furan L³³ had served as leads for development of antioxidant COX-2 inhibitors. Naturally occurring chromones such as stellatin derivatives³⁴, 2,6-Di-*tert*-butyl-4-thiomorpholin-4-




Figure 1. Modus operandi and biochemical relationship of anti-inflammatory and antioxidant agents in targeting radiation induced ROS formation.

ylmethylphenol hydrochloride³⁵ and crude plant extracts of *Tridax procumbis L.* and *Dioscorea opposite*³⁶ have also been reported to exert anti-inflammatory, antioxidant, and a variety of other pharmacological activities through inhibition of COX-2 mediated inflammatory responses³⁷⁻³⁸. The hypothesis with respect to the mechanism of action is described in Fig. 1.

6. COMPREHENSIVE DELIVERY SYSTEM FOR FEEDING

During times of high risk of radiation exposure, the person entering such area must ensure to bear the protective coverings in the form of suits, jacket, coat etc. The person who is basically operating in CBRN environment has to be satisfied of hunger and thirst. It becomes crucial to provide the suitable diet which not only takes care of basic nutrition, energy requirement but also enable the body's immunity to boost to fight against the potential threats. It keeps body hydrated and provides enough phytochemicals such as vitamins, antioxidants, and dietary fibers to overcome the detrimental effect of entities generated due to CBRN environment exposure³⁹. When the hands are held tight with the arms and ammunitions, the delivery system suitably fabricated/amalgamated with the suit can be very handy and user-friendly approach. The canister/bottle / pouch contains sterile food within must be capable of resisting

radiation exposure. It should be preferably a sterilizable and reusable system. The tubing mounting an orifice at one end opens in vicinity to the mouth cavity and can deliver item with a gentle press. The flow-able meal in liquid or semisolid form would be the choice of customer. A variety of menu could be designed and be delivered with above delivery system without introducing monotony in CBRN Meal. At present the design and development of such system is meager in most of developed and developing nations in general⁴⁰.

7. PROTECTION OF FOOD

Food may become contaminated from enemy employment of NBC weapons/agents or from terroristic contamination of food procurement facilities and food supplies. The NBC agents may be introduced during production or in the storage area of the procurement facility; while the product is in transit; at the military storage facility; or at the unit food service facility. Regardless of where the agent is used, the effect is the same; personnel will become ill if they consume the contaminated food. To ensure food safety, food must be inspected and monitored from its procurement until it is issued to the consumer. Throughout the channels, all services (Army, Navy, Marine, and Air Force) logistic and food service personnel must take precautions to protect substances from hazardous contamination³⁹⁻⁴⁰.

Designated experts are involved in the detection and monitoring of CBRN contaminated raw and processed food materials inclusive of tinned, canned and packed rations; before use, they must inspect all food suspected of being contaminated with NBC agents. They provide advice on the decontamination of food to unit personnel owning the food, or personnel performing the food decontamination as shown in Fig. 2. Depending on the type of contamination and packaging, the food may be; Consumed without being decontaminated; Decontaminated and then consumed; Destroyed. Some items

may be held to allow time for natural decay of nuclear or chemical contamination before consumption. The commander, with advice from veterinary/food safety expert, makes the decision on the disposition of the food⁴⁰ as shown in Table 2.

8. EFFECT OF CBRN AGENTS ON THE QUALITY OF FOOD AND THE DECONTAMINATION PROTOCOLS

The chemical contamination of the processed foods supplied in impermeable containers such as cans, bottles etc. can be resolved by immersing in boiling water/ soap water/ spraying with decontaminating solution-2 (DS2) for 30 minutes followed by rinsing and aeration (Table 2). And if the food is not packed in impermeable material and suspected to be contaminated with chemical agents it must be discarded. For biological decontamination of the packed food supplies the impermeable containers are boiled for 15 min/washed with soap and water followed by immersing in disinfectant solutions such as chlorine, bleaching solution, 5 per cent sodium carbonate, HTH (calcium hypochlorite) solution, super tropical bleach (STB) solution, 2 per cent per acetic acid. and aeration of the containers⁴¹. For nuclear decontamination the supplies must be washed with soap water followed by rinsing or brushing and wiping the contamination from surface. If the food is exposed the portion infected can be trimmed. The chemical agents such as Mustard, N-Mustards, Arsenicals,

DO's	Don'ts
<ul style="list-style-type: none"> • Consume only covered food products. • Eat canned foods. • Consume milk, fruit and vegetables bought or picked before release of toxic material and stored indoors in sealed packaging. • Consume peel able fruits and vegetables such as banana, oranges, cabbage, etc. • Decontaminate food storage containers thoroughly before opening. 	<ul style="list-style-type: none"> • Avoid water from open wells/ponds. • Avoid consumption of contaminated crops and vegetables, water or milk. • Avoid drinking fresh milk or eating fruits and vegetables grown in the affected area. • Avoid consumption of flours stored in jute bags. • Avoid dairy and meat products to avoid secondary contamination.

Figure 2. Dos and Don'ts for food consumption during CBRN scenario.

Table 2. Contagion and de-contagion methods³⁹⁻⁴²

	Avoid contamination-	Detection and Decontamination-
Priorities during CBRN scenarios	1- By using barriers 2- Specific entry and exit procedures 3- Preventing secondary contamination to spread	By specific methods (preferably remote sensing). Rapid methods must be practiced so as to sustain operation. Priorities the food group and packaging material.
Biological Toxins and microorganisms, biological derived agents	Mostly such contaminations do not change its physical appearance. Packaging material such as flexible films/ metal cans/ glass bottles etc. are highly impermeable for most of the biological threatening agent. Products handled in sealed storage containers such as freezers, refrigerated trucks, or rail cars are safe with respect to biological contamination till these are sealed or not exposed. Packing material is non-nutritive for microorganisms and therefore it would be non-life supportive, hence is self-decontaminating except for spores.	Rapid detection methods are essential to be applied for effective countermeasures at designated facility. Use boiling water/ Soap water for washing, use disinfectant such as bleaching powder, 5% Sodium carbonate solution, 2% per-acetic acid solution etc.
Chemical	Nerve agents / sulphur mustard, vesicants and arsenicals are most hazards and are either adsorbed or absorbed. In general causes gastrointestinal irritation/ systemic poisoning etc. Phosgene/hydrocyanic acid/ chemical irritants and toxic smokes are main culprit for food contamination not protected suitably. Fatty foods, water rich food such as fresh fruits and veg's having crystalline texture and grainy foods (sugar/ grains) attract deep penetration of nerve agents.	Quick tests kits based on paper strips/ handy colorimeter are extremely helpful for rapid detection and to suitable decide the next course of action in contaminated to sustain the fight. Decontamination is rather challenging but Boil water immersion, spaying disinfectants solution may help up to certain extent.
Radio nuclear	Direct contamination results when the reagents fallout on food and leads to its surface contamination Washing with disinfectant solution/ Brushing the infected surface/ wiping chlorine/ trimming of contaminated part in case unpacked food	Indirect contamination is by eating the contaminated foods Induced contamination happens when Exposed to sufficient neutron flux

nerve agents, Phosgene, Cyanogen agents, Irritants, smoke and white phosphorus imparts acidic and bitter taste, discoloration of the natural pigment and objectionable flavour to the foods exposed⁴². Usually the sealed metal cans, airtight glass bottles followed by metal foil laminates, wax papers, multilayer bags, cellophane, card board, wooden crates provide good protection from chemical vapors, and liquid chemical hazards of CBRN Environment, but the paper, canvas and textiles provide no protection to the food. The liquid chemical agents basically show their persistency in varied climatic conditions. They last long in calm sunny, snow bound areas (-10 °C) followed by sunny around (20 °C) with light breeze while wet and heavy wind do not favor their persistency⁴³.

9. CONCLUSIONS

The future war conditions and patterns are not obvious and the use of chemical, biological, radio-nuclear agent will be in any form may lead to serious detrimental effect on present living being in the area of incident. In such conditions protecting life with suitable life support aids would be an effective step to cope and overcome the current challenges of the war field. At times the food becomes a matter of survival and strength against fighting these situations and provides required source of energy and vital nutrients which can protect or delay the harmful effect of CBRN entities. At DRDO-Defence Food Research Laboratory the processed food are being packed in multilayer flexible packing material subsequently by secondary and tertiary packing material. The laboratory also has experience and expertise in the development of packaging materials for protective packaging of foods during NBC conditions. These materials act as strong barrier however with availability of infrastructure and expertise in various unit process operations which can be used in unison in furtherance of multi-target, combination processing and preservation, focused work for the development of nutrient supplement, its delivery system and packing which can address various challenges present in the CBRN conditions to survive life and to sustain missions is essential for future war scenario.

REFERENCES

1. Cameron, G. & Pate, J. Covert biological weapons attacks against agricultural targets: Assessing the impact against US agriculture. *Terrorism Political Violence*, 2001, **13**(3), 61-82.
doi: 10.1080/09546550109609690.
2. Carus, W.S. Rajneeshees. Toxic terror: Assessing terrorist use of chemical and biological weapons. In *Encyclopedia Bioterrorism Defefnce*, edited by Tucker JB. Cambridge, MA: MIT Press, 2000. 115–137.
3. Carus, W.S. Bioterrorism and biocrimes: The illicit use of biological agents since 1900. *The Minerva Group, Inc.*, 2002.
4. Prise, K.M.; Schettino, G.; Folkard, M. & Held, K.D. New insights on cell death from radiation exposure. *The Lancet Oncol.*, 2005, **6**(7), 520-528.
doi: 10.1016/S1470-2045(05)70246-1.
5. Maurya, D.K.; Devasagayam, T. & Nair, C.K.K. Some novel approaches for radioprotection and the beneficial effect of natural products, 2006.
6. Samarth, R.M.; Samarth, M. & Matsumoto, Y. Utilisation of cytogenetic biomarkers as a tool for assessment of radiation injury and evaluation of radiomodulatory effects of various medicinal plants– A review. *Drug Des., Dev. Ther.*, 2015, **9**, 5355.
doi: 10.2147/DDDT.S91299.
7. Fischer, N.; Seo, E.J. & Efferth, T. Prevention from radiation damage by natural products. *Phytomedicine*, 2017.
doi: 10.1016/j.phymed.2017.11.005
8. Smina, T.P.; Maurya, D.K.; Devasagayam, T.P.A. & Janardhanan, K.K. Protection of radiation induced DNA and membrane damages by total triterpenes isolated from *Ganoderma lucidum* (Fr.) P. Karst. *Chem.-Biol. Interact.*, 2015, **233**, 1-7.
doi: 10.1016/j.cbi.2015.03.019.
9. Painuli, S. & Kumar, N. Prospects in the development of natural radioprotective therapeutics with anti-cancer properties from the plants of Uttarakhand region of India. *J. Ayurveda Integr. Med.*, 2016, **7**(1), 62-68.
doi: 10.1016/j.jaim.2015.09.001.
10. Laqueur, W. No end to war: Terrorism in the twenty-first century. Bloomsbury Publishing, 2003.
11. Miller, J.; Broad, W.J. & Engelberg, S. Germs: Biological weapons and America's secret war. *Simon and Schuster*, 2012.
12. Jang, M.; Kim, H.K.; Choi, C.W. & Kang, C.S. Age-dependent potassium iodide effect on the thyroid irradiation by ¹³¹I and ¹³³I in the nuclear emergency. *Radiat. Prot. Dosim.*, 2008, **130**(4), 499-502.
doi: 10.1093/rpd/ncn068.
13. Krupnick, A.; Taylor, M.; Batz, M.; Hoffmann, S.; Tick, J.; Morris, G. & Sherman, D. Identifying the most significant microbiological foodborne hazards to public health: A new risk ranking model (*No. dp-frsc-dp-01*), 2004.
14. De la Roche, H.M.; Seagrove, S.; Mehta, A.; Divekar, P.; Campbell, S. & Curnow, A. Using natural dietary sources of antioxidants to protect against ultraviolet and visible radiation-induced DNA damage: An investigation of human green tea ingestion. *J. Photochem. Photobiol. B: Biol.*, 2010, **101**(2), 169-173.
15. Pate, J.; Ackerman, G. & McCloud, K. WMD terrorism chronology: Incidents involving sub-national actors and chemical, biological, radiological, or nuclear materials. Center for Nonproliferation Studies, Monterey Institute for International Studies, 2000, Available at <http://cns.miis.edu/pubs/reports/cbrn2k.htm>. (Accessed on 20 Nov 2017)
16. Turnbull, W. & Abhayaratne, P. WMD terrorism chronology: Incidents involving sub-national actors and chemical, biological, radiological, and nuclear materials. Monterey Institute of International Studies: Center for Nonproliferation Studies, Washington, DC, 2002.
17. Devipriya, N.; Sudheer, A.R.; Srinivasan, M. & Menon, V.P. Quercetin ameliorates gamma radiation-induced DNA damage and biochemical changes in human peripheral blood lymphocytes. *Mutat. Res. Genet. Toxicol. Environ.*

- Mutagen.*, 2008, **654**(1), 1-7.
doi: 10.1016/j.mrgentox.2008.03.003.
18. Maurya, Dharmendra K.; Devasagayam, Thomas & Nair, Cherupally Krishnan K. Some novel approaches for radioprotection and the beneficial effect of natural products, 2006.
 19. Smith, Tyler A.; Daniel R. Kirkpatrick; Sean Smith; Trevor, K. Smith; Pearson, Tate; Kailasam, Aparna; Kortney, Z.; Herrmann, Johanna Schubert & Agrawal, Devendra K. Radioprotective agents to prevent cellular damage due to ionising radiation. *J. Transl. Med.*, 15, 2017, **232**.
 20. Whitaker, T.B. & Johansson, A.S. Sampling uncertainties for the detection of chemical agents in complex food matrices. *J. Food Prot.*, 2005, **68**(6), 1306-1313.
doi: 10.4315/0362-028X-68.6.1306.
 21. Weiss, J.F. & Landauer, M.R. Radioprotection by antioxidants. *Ann. N.Y. Acad. Sci.* 2000, **899**(1), 44-60.
doi: 10.1186/s12967-017-1338-x.
 22. Weiss, J.F. & Landauer, M.R. Protection against ionising radiation by antioxidant nutrients and phytochemicals. *Toxicology*, 2003, **189**(1), 1-20.
doi: 10.1016/S0300-483X(03)00149-5.
 23. Riley, P.A. Free radicals in biology: oxidative stress and the effects of ionising radiation. *Int. J. Radiat. Biol.*, 1994, **65**(1), 27-33.
 24. Wambi, C.O.; Sanzari, J.K., Sayers, C.M., Nuth, M.; Zhou, Z.; Davis, J. & Kennedy, A.R. Protective effects of dietary antioxidants on proton total-body irradiation-mediated hematopoietic cell and animal survival. *Radiat. Res.*, 2009, **172**(2), 175-186.
doi: 10.1667/RR1708.1.
 25. Slade, D. & Radman, M. Oxidative stress resistance in *Deinococcus radiodurans*. *Microbiol. Mol. Biol. Rev.*, 2011, **75**(1), 133-191.
doi: 10.1128/MMBR.00015-10.
 26. Sprando, R.L.; Collins, T.F.X.; Black, T.; Olejnik, N.; Ramos-Valle, M. & Ruggles, D. Acute toxicity of sodium arsenite in a complex food matrix. *Food Chem. Toxicol.*, 2007, **45**(9), 1606-1613.
doi: 10.1016/j.fct.2007.02.026.
 27. Shukla, S.K.; Chaudhary, P.; Kumar, I.P.; Samanta, N.; Afrin, F.; Gupta, M.L. & Sharma, R.K. Protection from radiation-induced mitochondrial and genomic DNA damage by an extract of *Hippophae rhamnoides*. *Environ. Mol. Mutagen.*, 2006, **47**(9), 647-656.
doi: 10.1002/em.20251.
 28. Bengmark, S. Curcumin, An atoxic antioxidant and natural NFκB, cyclooxygenase-2, lipooxygenase, and inducible nitric oxide synthase inhibitor: A shield against acute and chronic diseases. *J. Parenter. Enteral Nutr.*, 2006, **30**(1), 45-51.
doi: 10.1177/014860710603000145.
 29. Bandgar, B.P.; Kinkar, S.N.; Chavan, H.V.; Jalde, S.S.; Shaikh, R.U. & Gacche, R.N. Synthesis and biological evaluation of asymmetric indole curcumin analogs as potential anti-inflammatory and antioxidant agents. *J. Enzyme Inhib. Med. Chem.*, 2014, **29**(1), 7-11.
doi: 10.3109/14756366.2012.743536.
 30. Candelario-Jalil, E.; de Oliveira, A.C.P.; Gräf, S.; Bhatia, H.S.; Hüll, M.; Muñoz, E. & Fiebich, B.L. Resveratrol potently reduces prostaglandin E 2 production and free radical formation in lipopolysaccharide-activated primary rat microglia. *J. Neuroinflammation*, 2007, **4**(1), 25.
doi: 10.1186/1742-2094-4-25.
 31. Ahmad, S.; Israf, D.A.; Lajis, N.H.; Shaari, K.; Mohamed, H.; Wahab, A.A.; Ariffin K.T.; Hoo, W.Y.; Aziz, N.A.; Kadir, A.A.; Sulaiman, M.R. & Somchit M.N. Cardamonin, inhibits pro-inflammatory mediators in activated RAW 264.7 cells and whole blood. *Eur. J. Pharmacol.*, 2006, **538**(1-3), 188-194.
doi: 10.1016/j.ejphar.2006.03.070.
 32. Seo, K.; Yang, J.H.; Kim, S.C.; Ku, S.K.; Ki, S.H. & Shin, S.M. The antioxidant effects of isorhamnetin contribute to inhibit COX-2 expression in response to inflammation: A potential role of HO-1. *Inflammation*, 2014, **37**(3), 712-722.
doi: 10.1007/s10753-013-9789-6.
 33. Nishio, K.; Fukuhara, A.; Omata, Y.; Saito, Y.; Yamaguchi, S.; Kato, H.; Yoshida, Y. & Niki, E. Characterisation of novel furan compounds on the basis of their radical scavenging activity and cytoprotective effects against glutamate-and lipopolysaccharide-induced insults. *Bioorg. Med. Chem.*, 2008, **16**(24), 10332-10337.
 34. Gautam, R.; Jachak, S.M.; Kumar, V. & Mohan, C.G. Synthesis, biological evaluation and molecular docking studies of stellatin derivatives as cyclooxygenase (COX-1, COX-2) inhibitors and anti-inflammatory agents. *Bioorg. Med. Chem. Lett.*, 2011, **21**(6), 1612-1616.
doi: 10.1016/j.bmc.2008.10.038.
 35. Ziakas, G.N.; Rekkas, E.A.; Gavalas, A.M.; Eleftheriou, P.T. & Kourounakis, P.N. New analogues of butylated hydroxytoluene as anti-inflammatory and antioxidant agents. *Bioorg. Med. Chem.*, 2006, **14**(16), 5616-5624.
doi: 10.1016/j.bmc.2006.04.030.
 36. Yang, M.H.; Yoon, K.D.; Chin, Y.W.; Park, J.H. & Kim, J. Phenolic compounds with radical scavenging and cyclooxygenase-2 (COX-2) inhibitory activities from *Dioscorea opposita*. *Bioorg. Med. Chem.*, 2009, **17**(7), 2689-2694.
doi: 10.1016/j.bmc.2009.02.057.
 37. Hirata, A.; Murakami, Y.; Shoji, M.; Kadoma, Y. & Fujisawa, S. Kinetics of radical-scavenging activity of hesperetin and hesperidin and their inhibitory activity on COX-2 expression. *Anticancer Res.*, 2005, **25**(5), 3367-3374.
 38. Laube, M.; Kniess, T. & Pietzsch, J. Development of antioxidant COX-2 inhibitors as radioprotective agents for radiation therapy—A hypothesis-driven review. *Antioxidants*, 2016, **5**(2), 14.
doi: 10.3390/antiox5020014.
 39. Weiss, J.F.; & Landauer, M.R. History and development of radiation-protective agents. *Int. J. Radiat. Biol.*, 2009, **85**(7), 539-573.
doi: 10.1080/09553000902985144.
 40. Thompson, J.S.; Chu, Y.; Glass, J.; Tapp, A.A. & Brown, S.A. The manganese superoxide dismutase mimetic,

M40403, protects adult mice from lethal total body irradiation. *Free Radical Res.*, 2010, **44**(5), 529-540.

doi: 10.3109/10715761003649578.

41. Velioglu-Ögünç, A.; Şehirli, Ö.; Toklu, H.Z.; Özyurt, H.; Mayadağlı, A.; Ekşioğlu-Demiralp, E. & Şener, G. Resveratrol protects against irradiation-induced hepatic and ileal damage via its anti-oxidative activity. *Free Radical Res.*, 2009, **43**(11), 1060-1071.
doi: 10.1080/10715760903171100.
42. Kumar, M.; Sharma, M.K.; Saxena, P.S. & Kumar, A. Radioprotective effect of Panax ginseng on the phosphatases and lipid peroxidation level in testes of Swiss albino mice. *Biol. Pharm. Bull.*, 2003, **26**(3), 308-312.
doi: 10.1248/bpb.26.308.
43. Devi, P. & Ganasoundari, A. Modulation of glutathione and antioxidant enzymes by *Ocimum sanctum* and its role in protection against radiation injury. *Indian J. Exp. Biol.*, 1999, **37**(3), 262-268.
44. Dreger, S.; Pfänder, M.; Christianson, L.; Lhachimi, S. K. & Zeeb, H. The effects of iodine blocking following nuclear accidents on thyroid cancer, hypothyroidism, and benign thyroid nodules: Design of a systematic review. *Syst. Rev.*, 2015, **4**(1), 126.
doi: 10.1186/s13643-015-0106-3.
45. Gee, C.A.; Kittridge, K.J. & Willson, R.L. Peroxy free radicals, enzymes and radiation damage: Sensitisation by oxygen and protection by superoxide dismutase and antioxidants. *Br. J. Radiol.*, 1985, **58**(687), 251-256.
doi: 10.1259/0007-1285-58-687-251.
46. Nambiar, D.; Rajamani, P. & Singh, R.P. Effects of phytochemicals on ionisation radiation-mediated carcinogenesis and cancer therapy. *Mutat. Res. Rev. Mutat. Res.*, 2011, **728**(3), 139-157.
doi: 10.1016/j.mrrev.2011.07.005.
47. Belay, A.; Ota, Y.; Miyakawa, K. & Shimamatsu, H. Current knowledge on potential health benefits of Spirulina. *J. Appl. Phycol.*, 1993, **5**(2), 235-241.

ACKNOWLEDGEMENT

The authors are grateful and acknowledge the constructive suggestions, support, and meticulous guidance of Head Grain Science & Technology Division and Head Fruits & Vegetable Technology Division.

CONTRIBUTORS

Mr. Dev Kumar Yadav has received his MSc (Food Technology) from University of Allahabad, Allahabad and currently pursuing his PhD (Food Science) from Bharathiar University. He is presently working as Scientist 'D' at DRDO-Defence Food Research Laboratory, Mysore. He has published more than 20 publications in journals. His research areas includes modification of starches for specific food applications, development of microwave based disinfection process, instant foods based on cereals, pulses & Millets, development of imitated milk products from decorticated sesame seeds, design and development of ration storage system for Service specific applications. He is the lead writer of the present paper.

Mrs Janifer Raj Xavier obtained her MSc (Agricultural Biotechnology) from Tamil Nadu Agricultural University, Coimbatore and currently pursuing her PhD (Biotechnology) from Bharathiar University. Presently working as Scientist 'D' at DRDO-Defence Food Research Laboratory, Mysore. She had published 35 research paper in journals. She has given inputs related to modus operandi and biochemical relationship of anti-inflammatory and antioxidant agents in targeting radiation induced ROS formation radio-protected herbs and contributed in writing of manuscript.

Dr OP Chauhan has received his PhD (Food Technology) from GB Pant University, Pantnagar, in 2002. Presently working as Scientist E at DRDO-Defence Food Research Laboratory, Mysore. He has 75 research publications, 11 book chapters, 1 book, and 9 patent to his credit. He has vast experience in processing of various value added products from fruits and vegetable as well as extension of their shelf life. He has meticulously contributed in the manuscript preparation.

Mr Prakash Eknath Patki is working as Scientist G and Head of Division in Fruits and Vegetable Technology at DRDO-Defence Food Research Laboratory, Mysore. He has worked in the development of instant foods based on cereals, pulses and millets. He has published 40 research papers and 10 patents to his credit. He has contributed meticulously for the suitable content and layout of the manuscript.

Dr RK Sharma received his M Pharm (Pharmaceutical Chemistry) from Panjab University and PhD from University of Delhi. He was former Director of DRDO-Defence Food Research Laboratory, Mysore. He has made significant contributions in new drugs, novel drugs delivery systems, herbal radioprotectors, herbal biothreat mitigators, and nutraceuticals. He has contributed meticulously for the suitable content and layout of the manuscript.